

# euromedica hannover 2012

.....

31. MAI — 01. JUNI 2012

## INTERNATIONALER MEDIZINISCHER KONGRESS

### Programm Abstracts

Europäische Akademie für Naturwissenschaften, e.V. Hannover  
Europäische Wissenschaftliche Gesellschaft e.V Hannover

Vorbereitender Ausschuss:

G. Tyminskiy (Hanover, Germany)  
W. Fischer (Berlin, Germany)  
L. Royt (Hanover, Germany)  
V. Astashov (Moscow, Russia)

V. Geldt (Moscow, Russia)  
K. Togusbaeva (Almaty, Kazakhstan)  
T. Abdufataev (Dushanbe, Tajikistan)  
V. Sokolov (Odessa, Ukraine)

ISSN 2194-3524

Herausgeber:

Dr. med. G. Tyminskiy  
Europäische Wissenschaftliche Gesellschaft  
Sutelstr. 50A  
30659 Hannover  
Tel. +49 (0)511 390 - 8088  
[www.congress-euromedica.de](http://www.congress-euromedica.de)  
E-mail: [ewg2002@googlemail.com](mailto:ewg2002@googlemail.com)

were analyzed by immunoenzyme automated method with apparatus «Emmulite».

Results. In 12-14 and 28-30 days after beginning of treatment there was a positive dynamic of most clinic and laboratory parameters: decrease of dyspnea from  $2,74 \pm 0,04$  to  $2,21 \pm 0,04$  at 12-14<sup>th</sup> days and to  $2,2 \pm 0,15$  at 28-30<sup>th</sup> days ( $p < 0,05$ ), patients noticed significant improvement of cough at 28-30<sup>th</sup> days ( $p < 0,05$ ), towards the end of period of treatment intensity of such symptoms as heartburn (47%), eructation (21%), dysphagia (9,8%) and it's connection with changing of pose (29,5%) was decreased ( $p < 0,05$ ). At 28-30<sup>th</sup> days of treatment significant increase of cortisol, testosterone and estradiol level in blood was detected. Evident increase of vegetative reactivity (SDNN) in 46,7% ( $p = 0,0002$ ) and decrease of parasympathetic activity (rMSSD) in 31,8% ( $p < 0,05$ ) occurred. Beneficial dynamic of test indices had continued after 6 and 12 months. Invalid dynamic of clinic and laboratory data was in placebo group. In 3<sup>th</sup> group there weren't significant changes of tested indices.

Conclusion. Experimentation has shown that combination of basic pharmacotherapy and repeated courses of laser therapy has the best efficiency and the most evident beneficial effect on clinical course of BA associated with GERD because of alternate action and activation of adaptive and compensatory response. This method of treatment influences on different parts of pathogenesis of BA and GERD. Consequently, introduced method of treatment makes it possible to increase effectiveness of therapy, shorten patient's stay in hospital, corrects level of hormones and vegetative imbalance as well as reduces drugs dose.

K.V. Nikitin

**THE TREATMENT OF PATIENTS WITH HERNIAS  
OF INTERVERTEBRAL DISKS WITH THE HELP OF SCENAR-  
THERAPY**

*Medical and Diagnostic Center, Tomsk, Russia*

The difficulty of pathology at hernias of intervertebral disks is well-known. It is connected with serious violation of exchange processes and is accompanied by heavy symptomatology from a pain syndrome to the full impossibility to go.

The treatment is interfaced to great difficulties. In orthodox medicine it is dispersed to the conflict against symptoms or operative intervention which is, in fact, palliative action as doesn't change factors brought to degenerate processes. However, it is not a simple task for the skenar-therapist as to restore an exchange for a long time in separately taken site is not possible, it is necessary to restore it in all an organism, and it, almost always, a problem.

We chose skenar-therapy as the main method of treatment because it allows to restore various systems of an organism at their damage. Thanks to some features (existence of the express diagnostics, return biological communication, influence by neurosimilar impulses), skenar-therapy acts as a universal regulator and the adjuster of physiological processes.

## Program Abstracts

We treated 186 people. Age – from 24 to 68 years. Number of treatment courses – from 5 to 11. Duration of a course – from 15 to 25 days. Duration of all treatment from 5 months to 2 years. In all cases the existence of hernia of one or several intervertebral disks of a backbone is confirmed by means of tomographic research.

At receipt complaints to pains in various parts of a backbone, in buttocks, in feet, a dumbness of the top and bottom extremities, weakness, a hyper tone and an undernourishment of muscles can be possible. In the anamnesis numerous inflammatory processes (tonsillitis, pharyngitises, otitises, antritises) at which treatment applied massive doses of antibiotics, and since the childhood. 34 (~18 %) patients are diagnosed stomach ulcer of a stomach and a duodenal gut, 86 (~46 %) patients had the phenomena enterocolitis and cholecystitis, 22 (~12 %) – hepatitis B. 120 (~64 %) patients had a considerable excess of weight and violation of a fatty exchange.

From the first sessions we explained to our patients that their real condition was the result of long combined influence of damaging factors of environment in which they lived and, partly, created. We analyzed a dietary habits in details and introduced all necessary amendments.

Steady positive result in the form of disappearance of pains, improvements of the general condition, restoration of working capacity is reached at 129 (~69 %) patients (follow-up from 2 to 9 years). From this number 96 (~51 %) patients by tomographic research had reduction of the phenomena of osteochondrosis, restoration of the damaged fibrous rings and disappearance of hernias of intervertebral disks.

From the remained 57 patients 19, having got rid of pains after the first plan of treatment, didn't appear any more and their destiny isn't known to us. The other 38 didn't want to change something in their life, especially to starve. After two courses of treatment they received considerable improvement in the form of disappearance of pains. Now they continue to visit us two-three times a year in process of renewal of painful feelings.

Thus, the treatment of similar diseases demands a complex influence and depends not only on qualification of the doctor and correctness of the chosen technique, but also on the courage of determination and understanding of the patient. And skenar-therapy can be considered as rather adequate method of treatment of similar conditions.

T.V. Novikova  
Y.S. Pekker  
O.V. Vorobeichikova

### COMPETENCY OF A CYBERNETICS DOCTOR AS A CONTRIBUTION TO MEDICAL WORK

*The Siberian State Medical University, Tomsk, Russia*

The qualification of a cybernetics doctor is given to a graduate of a medical university in accordance with the Russian Federal Standard of Education. The competency is seen as the ability to demonstrate knowledge and skills, which have been acquired according to the qualification characteristics, in practice.